

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for **spla2 inhibitor and vascular** Preview Go

Clear

About Entrez  
Text Version

Limits Preview/Index **History** Clipboard Details

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy

E-Utilities

- Search History will be lost after eight hours of inactivity.
- Search numbers may not be continuous; all searches are represented.
- To save search indefinitely, click query # and select Save in My NCBI.
- To combine searches use #search, e.g., #2 AND #3 or click query # for more options.

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Search	Most Recent Queries	Time	Result
#20	Search <b>spla2 inhibitor and vascular</b>	16:12:54	<u>18</u>
#19	Search <b>spla2 inhibitor and amyloid</b>	16:12:19	<u>1</u>
#2	Search <b>spla2 inhibitor and hypertension</b>	16:12:12	<u>4</u>
#18	Search <b>spla2 inhibitor and vasoactive</b>	16:12:05	<u>0</u>
#1	Search <b>spla2 inhibitor</b>	16:11:56	<u>205</u>
#10	Search <b>beta amyloid and hypertension</b>	16:08:45	<u>90</u>
#4	Search <b>spla2 and hypertension</b>	15:57:37	<u>4</u>
#9	Search <b>amyloid and hypertension</b>	15:02:13	<u>401</u>
#7	Search <b>secretory phospholipase A2 and alzheimers</b>	13:55:45	<u>2</u>
#6	Search <b>secretory phospholipase A2</b>	13:55:37	<u>968</u>
#5	Search <b>secretory phospholipase A2 and hypertension</b>	13:55:25	<u>4</u>
#3	Search <b>spla2 inhibitor and vasospasm</b>	13:54:41	<u>0</u>

Clear History

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

## EAST Search History

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	65	"5700816"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 11:21
L2	56	"spla2 inhibitor"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 11:22
L3	8	"spla2 inhibitor" and "alzheimer"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 11:24
L4	0	secretory adj phospholipase adj inhibitor	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 11:47
L5	0	"secretory phospholipase inhibitor"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 11:47
L6	8	secretory near phospholipase near inhibitor	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 11:48
L7	65	"5654326"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 12:27
L8	1	"spla2 inhibitor" and "hypertension"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 12:50

## EAST Search History

L9	0	"spla2 inhibitor" and "vascular disease"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 13:53
L10	0	"spla2 inhibitor" and "vasospasm"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 13:53
L11	10	"6,166,017"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 14:11
L12	4	"6642236"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 14:11
L13	14	"cortisol inhibitor"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 14:12
L14	3	"6642236" and "cortisol inhibitor"	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 14:12
L15	0	oleyloxyethylphosphocholine	US-PGPUB; USPAT; USOCR; FPRS; EPO; JPO; DERWENT	OR	OFF	2008/01/25 16:43

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTAJDA1614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	AUG 06	CAS REGISTRY enhanced with new experimental property tags
NEWS	3	AUG 06	FSTA enhanced with new thesaurus edition
NEWS	4	AUG 13	CA/CAPplus enhanced with additional kind codes for granted patents
NEWS	5	AUG 20	CA/CAPplus enhanced with CAS indexing in pre-1907 records
NEWS	6	AUG 27	Full-text patent databases enhanced with predefined patent family display formats from INPADOCDB
NEWS	7	AUG 27	USPATOLD now available on STN
NEWS	8	AUG 28	CAS REGISTRY enhanced with additional experimental spectral property data
NEWS	9	SEP 07	STN AnaVist, Version 2.0, now available with Derwent World Patents Index
NEWS	10	SEP 13	FORIS renamed to SOFIS
NEWS	11	SEP 13	INPADOCDB enhanced with monthly SDI frequency
NEWS	12	SEP 17	CA/CAPplus enhanced with printed CA page images from 1967-1998
NEWS	13	SEP 17	Caplus coverage extended to include traditional medicine patents
NEWS	14	SEP 24	EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS	15	OCT 02	CA/CAPplus enhanced with pre-1907 records from Chemisches Zentralblatt
NEWS	16	OCT 19	BEILSTEIN updated with new compounds
NEWS	17	NOV 15	Derwent Indian patent publication number format enhanced
NEWS	18	NOV 19	WPIX enhanced with XML display format
NEWS	19	NOV 30	ICSD reloaded with enhancements
NEWS	20	DEC 04	LINPADOCDB now available on STN
NEWS	21	DEC 14	BEILSTEIN pricing structure to change
NEWS	22	DEC 17	USPATOLD added to additional database clusters
NEWS	23	DEC 17	IMSDRUGCONF removed from database clusters and STN
NEWS	24	DEC 17	DGENE now includes more than 10 million sequences
NEWS	25	DEC 17	TOXCENTER enhanced with 2008 MeSH vocabulary in MEDLINE segment
NEWS	26	DEC 17	MEDLINE and LMEDLINE updated with 2008 MeSH vocabulary
NEWS	27	DEC 17	CA/CAPplus enhanced with new custom IPC display formats
NEWS	28	DEC 17	STN Viewer enhanced with full-text patent content from USPATOLD
NEWS	29	JAN 02	STN pricing information for 2008 now available
NEWS	30	JAN 16	CAS patent coverage enhanced to include exemplified prophetic substances
NEWS EXPRESS	19	SEPTEMBER 2007:	CURRENT WINDOWS VERSION IS V8.2, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 19 SEPTEMBER 2007.
NEWS HOURS	STN Operating Hours Plus Help Desk Availability		
NEWS LOGIN	Welcome Banner and News Items		
NEWS IPC8	For general information regarding STN implementation of IPC 8		

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 16:45:32 ON 25 JAN 2008

=> file registry

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 16:45:44 ON 25 JAN 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 24 JAN 2008 HIGHEST RN 1000773-19-2

DICTIONARY FILE UPDATES: 24 JAN 2008 HIGHEST RN 1000773-19-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=> E "OLEYLOXYETHYLPHOSPHOCHOLINE"/CN 25

E1 1 OLEYLONITRILE/CN

E2 1 OLEYLOXYETHYL CIDOFOVIR/CN

E3 0 --> OLEYLOXYETHYLPHOSPHOCHOLINE/CN

E4 1 OLEYLOXYPROPYL-N,N-DIMETHYLAMINE/CN

E5 1 OLEYLPALMITAMIDE/CN

E6 1 OLEYLPHENOL/CN

E7 1 OLEYLPHOSPHORYLETHANOLAMINE/CN

E8 1 OLEYLPROPYLENEDIAMINE/CN

E9 1 OLEYLSARCOSINE N-HEPTADECYL-1,3-PROPANEDIAMINE SALT/CN

E10 1 OLEYLSARCOSINE SODIUM SALT/CN

E11 1 OLEYLSTEARYLAMINE/CN

E12 1 OLEYLSUCCINIC ACID/CN

E13 1 OLEYLSUCCINIC ANHYDRIDE/CN

E14 1 OLEYLTRIMETHYLAMMONIUM BROMIDE/CN

E15 1 OLEYLTRIMETHYLAMMONIUM CHLORIDE/CN

E16 1 OLEYLTRIMETHYLENEDIAMINE/CN

E17 1 OLEYLTRIOCTADECYLAMMONIUM BROMIDE/CN

E18 1 OLF-1/EBF TRANSCRIPTION FACTOR (CAENORHABDITIS ELEGANS STRAIN N2

GENE UNC-3)/CN


E19 1 OLF-1/EBF-LIKE-1(8) TRANSCRIPTION FACTOR (MOUSE STRAIN CD-1  
8-AMINO ACID INSERT ISOFORM)/CN

E20 1 OLF-1/EBF-LIKE-2 (9L) TRANSCRIPTION FACTOR (MOUSE STRAIN CD-1  
 LONG ISOFORM 9L)/CN  
 E21 1 OLF-1/EBF-LIKE-2 (OS) TRANSCRIPTION FACTOR (MOUSE STRAIN CD-1  
 SHORT ISOFORM 03)/CN  
 E22 1 OLF-1/EBF-LIKE-3 TRANSCRIPTION FACTOR (MOUSE STRAIN CD-1)/CN  
 E23 1 OLFACTOMEDIN (HUMAN CLONE DE10316701-SEQID-623 GENE OLFM1  
 ISOFORM 1)/CN  
 E24 1 OLFACTOMEDIN (RANA CATESBEIANA PRECURSOR REDUCED)/CN  
 E25 1 OLFACTOMEDIN 1 (HUMAN GENE OLFM1)/CN  
  
 => E "OOEPC"/CN 25  
 E1 1 OOE/CN  
 E2 1 OOENL/CN  
 E3 0 --> OOEPC/CN  
 E4 1 OOG4 PROTEIN (MOUSE STRAIN C57BL/6J CLONE IMAGE:30053268 GENE  
 OOG4)/CN  
 E5 1 OOGENESIN (MOUSE CLONE C0103)/CN  
 E6 1 OOGENESIN 2 (MOUSE STRAIN C57BL/6J CLONE MGC:60747  
 IMAGE:30050073)/CN  
 E7 1 OOGENESIN 3 (MOUSE STRAIN C57BL/6J CLONE MGC:60746  
 IMAGE:30049894)/CN  
 E8 1 OOGONIOL/CN  
 E9 1 OOGONIOL 1/CN  
 E10 1 OOGONIOL 2/CN  
 E11 1 OOGONIOL 3/CN  
 E12 1 OOGONIOL TETRAACETATE/CN  
 E13 1 OOH17N14M2/CN  
 E14 1 OOI/CN  
 E15 1 OOK 1/CN  
 E16 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM BERGHEI STRAIN  
 ANKA)/CN  
 E17 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM BERGHEI STRAIN  
 QIMR)/CN  
 E18 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM CHABAUDI ADAMI STRAIN  
 DS)/CN  
 E19 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM CHABAUDI ADAMI STRAIN  
 KA556)/CN  
 E20 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM CHABAUDI CHABAUDI  
 STRAIN AJ)/CN  
 E21 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM CHABAUDI CHABAUDI  
 STRAIN AS)/CN  
 E22 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM CHABAUDI CHABAUDI  
 STRAIN CB1455)/CN  
 E23 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM VINCKEI PETTERI )/CN  
 E24 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM VINCKEI VINCKEI )/CN  
 E25 1 OOKINETE SURFACE ANTIGEN PFS28 (PLASMODIUM YOELII NIGERIENSIS  
 )/CN

=> S 96720-06-8/RN  
 L1 1 96720-06-8/RN

=> DIS L1 1 SQIDE  
 THE ESTIMATED COST FOR THIS REQUEST IS 6.65 U.S. DOLLARS  
 DO YOU WANT TO CONTINUE WITH THIS REQUEST? (Y)/N:Y


L1 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2008 ACS on STN  
 RN 96720-06-8 REGISTRY  
 CN 3,5,8-Trioxa-4-phosphahexacos-17-en-1-aminium, 4-hydroxy-N,N,N-trimethyl-,  
 inner salt, 4-oxide (9CI) (CA INDEX NAME)  
 MF C25 H52 N O5 P  
 LC STN Files: BEILSTEIN\*, CA, CAPLUS, CASREACT, CHEMCATS, CSChem, MEDLINE,  
 TOXCENTER  
 (\*File contains numerically searchable property data)  
 DT.CA Caplus document type: Journal  
 RL.NP Roles from non-patents: PREP (Preparation)



1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> d 11

L1 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2008 ACS on STN  
RN 96720-06-8 REGISTRY  
ED Entered STN: 09 Jun 1985  
CN 3,5,8-Trioxa-4-phosphahexacos-17-en-1-aminium, 4-hydroxy-N,N,N-trimethyl-,  
inner salt, 4-oxide (9CI) (CA INDEX NAME)  
MF C25 H52 N O5 P  
LC STN Files: BEILSTEIN\*, CA, CAPLUS, CASREACT, CHEMCATS, CSCHEM, MEDLINE,  
TOXCENTER  
(\*File contains numerically searchable property data)



1 REFERENCES IN FILE CA (1907 TO DATE)  
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> s 11

L2 1 96720-06-8/RN

=> file medline caplus wpids uspatfull  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
7.22	7.43

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 16:49:58 ON 25 JAN 2008

FILE 'CAPLUS' ENTERED AT 16:49:58 ON 25 JAN 2008  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIDS' ENTERED AT 16:49:58 ON 25 JAN 2008  
COPYRIGHT (C) 2008 THE THOMSON CORPORATION

FILE 'USPATFULL' ENTERED AT 16:49:58 ON 25 JAN 2008  
CA INDEXING COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

=> s 11

'RN' IS NOT A VALID FIELD CODE

L3 8 L1

=> d 13 1-8 ibib, abs, hitstr

L3 ANSWER 1 OF 8 MEDLINE on STN

ACCESSION NUMBER: 2002725402 MEDLINE  
 DOCUMENT NUMBER: PubMed ID: 12489129  
 TITLE: Eicosanoids in insect immunity: bacterial infection stimulates hemocytic phospholipase A2 activity in tobacco hornworms.  
 AUTHOR: Tunaz Hasan; Park Youngjin; Buyukguzel Kemal; Bedick Jon C; Nor Aliza A R; Stanley David W  
 CORPORATE SOURCE: Insect Biochemical Physiology Laboratory, University of Nebraska, Lincoln 68583-0816, USA.  
 SOURCE: Archives of insect biochemistry and physiology, (2003 Jan) Vol. 52, No. 1, pp. 1-6.  
 Journal code: 8501752. ISSN: 0739-4462.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 (RESEARCH SUPPORT, NON-U.S. GOV'T)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200303  
 ENTRY DATE: Entered STN: 19 Dec 2002  
 Last Updated on STN: 4 Mar 2003  
 Entered Medline: 3 Mar 2003

AB Intracellular phospholipase A(2) (PLA(2)) is responsible for releasing arachidonic acid from cellular phospholipids, and is thought to be the first step in eicosanoid biosynthesis. Intracellular PLA(2)s have been characterized in fat body and hemocytes from tobacco hornworms, *Manduca sexta*. Here we show that bacterial challenge stimulated increased PLA(2) activity in isolated hemocyte preparations, relative to control hemocyte preparations that were challenged with water. The increased activity was detected as early as 15 s post-challenge and lasted for at least 1 h. The increased activity depended on a minimum bacterial challenge dose, and was inhibited in reactions conducted in the presence of oleyoxyethylphosphorylcholine, a site-specific PLA(2) inhibitor. In independent experiments with serum prepared from whole hemolymph, we found no PLA(2) activity was secreted into serum during the first 24 h following bacterial infection. We infer that a hemocytic intracellular PLA(2) activity is increased immediately an infection is detected. The significance of this enzyme lies in its role in launching the biosynthesis of eicosanoids, which mediate cellular immune reactions to bacterial infection.

Copyright 2003 Wiley-Liss, Inc.

L3 ANSWER 2 OF 8 MEDLINE on STN  
 ACCESSION NUMBER: 2002147899 MEDLINE  
 DOCUMENT NUMBER: PubMed ID: 11841807  
 TITLE: Identification of the phospholipase A(2) isoforms that contribute to arachidonic acid release in hypoxic endothelial cells: limits of phospholipase A(2) inhibitors.  
 AUTHOR: Michiels Carine; Renard Patricia; Bouaziz Najat; Heck Nathalie; Eliaers Francois; Ninane Noelle; Quarck Rozenn; Holvoet Paul; Raes Martine  
 CORPORATE SOURCE: Laboratoire de Biochimie et Biologie Cellulaire, Facultes Universitaires Notre Dame de la Paix, 61 rue de Bruxelles, 5000, Namur, Belgium.. carine.michiels@fundp.ac.be  
 SOURCE: Biochemical pharmacology, (2002 Jan 15) Vol. 63, No. 2, pp. 321-32.  
 Journal code: 0101032. ISSN: 0006-2952.  
 PUB. COUNTRY: England: United Kingdom  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 (RESEARCH SUPPORT, NON-U.S. GOV'T)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200203  
 ENTRY DATE: Entered STN: 8 Mar 2002  
 Last Updated on STN: 3 Apr 2002  
 Entered Medline: 27 Mar 2002



AB Changes in endothelium functions during ischemia are thought to be of importance in numerous pathological conditions, with, for instance, an increase in the release of inflammatory mediators like prostaglandins. Here, we showed that hypoxia increases phospholipase A(2) (PLA(2)) activity in human umbilical vein endothelial cells. Both basal PLA(2) activity and PG synthesis are sensitive to BEL and AACOCF3, respectively, inhibitors of calcium-independent PLA(2) (iPLA(2)) and cytosolic PLA(2) (cPLA(2)), while OPC, an inhibitor of soluble PLA(2) (sPLA(2)) only inhibited the hypoxia-induced AA release and PGF(2alpha) synthesis. Hypoxia does not alter expression of iPLA(2), sPLA(2) and cPLA(2) and cycloheximide did not inhibit PLA(2) activation, indicating that hypoxia-induced increase in PLA(2) activity is due to activation rather than induction. However, mRNA levels for sPLA(2) displayed a 2-fold increase after 2 hr incubation under hypoxia. BAPTA, an intracellular calcium chelator, partially inhibited the AA release in normoxia and in hypoxia. Direct assays of specific PLA(2) activity showed an increase in sPLA(2) activity but not in cPLA(2) activity after 2hr hypoxia. Taken together, these results indicate that the hypoxia-induced increase in PLA(2) activity is mostly due to the activation of sPLA(2).

L3 ANSWER 3 OF 8 MEDLINE on STN  
ACCESSION NUMBER: 2001301904 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 11226404  
TITLE: The involvement of phospholipase A(2) in ethanol-induced gastric muscle contraction.  
AUTHOR: Sim S S; Choi J C; Min D S; Rhie D J; Yoon S H; Hahn S J; Kim C J; Kim M S; Jo Y H  
CORPORATE SOURCE: Department of Pathophysiology, College of Pharmacy, Chung-Ang University, 221 Huksuk-dong, Dongjak-gu, Seoul 156-756, South Korea.  
SOURCE: European journal of pharmacology, (2001 Feb 16) Vol. 413, No. 2-3, pp. 281-5.  
Journal code: 1254354. ISSN: 0014-2999.  
PUB. COUNTRY: Netherlands  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200105  
ENTRY DATE: Entered STN: 4 Jun 2001  
Last Updated on STN: 4 Jun 2001  
Entered Medline: 31 May 2001

AB To understand the underlying mechanism of ethanol in tonic contraction, the effect of ethanol on phospholipase A(2) and phospholipase C activities and the effects of phospholipase inhibitors on ethanol-induced contraction of cat gastric smooth muscle were tested. Circular muscle strips (2.0 x 0.2 cm) obtained from the fundus of cat stomach were used to measure isometric contraction. Ethanol elicited tonic contraction and activated phospholipase A(2) activity in a dose-dependent manner. Phospholipase A(2) inhibitors, manoalide (0.1--10 microM) and oleyloxyethyl phosphorylcholine (1--10 microM), significantly inhibited ethanol-induced contraction. Furthermore, 342 mM ethanol-induced contraction was significantly inhibited by cyclooxygenase inhibitors, ibuprofen (10--100 microM) and indomethacin (10--100 microM), but not by lipoxygenase inhibitors. On the other hand, phospholipase C inhibitors had no effect on ethanol-induced contraction, indicating that phospholipase C is not involved in ethanol-induced contraction. It is suggested from the above results that ethanol-induced contraction in cat gastric smooth muscle is, in part, mediated by phospholipase A(2) and cyclooxygenase pathways.

L3 ANSWER 4 OF 8 MEDLINE on STN  
ACCESSION NUMBER: 2000106581 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 10643787  
TITLE: Novel strategies for opposing murine microglial activation.  
AUTHOR: Paris D; Town T; Mullan M

CORPORATE SOURCE: The Roskamp Institute, University of South Florida, Tampa  
33613, USA.. dparis@com1.med.usf.edu  
SOURCE: Neuroscience letters, (2000 Jan 7) Vol. 278, No. 1-2, pp.  
5-8.  
Journal code: 7600130. ISSN: 0304-3940.  
PUB. COUNTRY: Ireland  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200002  
ENTRY DATE: Entered STN: 9 Mar 2000  
Last Updated on STN: 9 Mar 2000  
Entered Medline: 22 Feb 2000

AB Pathologic microglial activation is believed to contribute to progressive neuronal damage in neurodegenerative diseases by the release of potentially neurotoxic agents, such as pro-inflammatory cytokines including tumor necrosis factor alpha (TNF-alpha). Using cultured N9 microglial cells, we have examined the regulation of TNF-alpha following endotoxic insult with lipopolysaccharide (LPS), focusing on the role of the pro-inflammatory phospholipase A2/mitogen activated protein kinase/arachidonic acid/cyclo-oxygenase-2 cascade and the nitric oxide/cGMP pathway. Data show that various inhibitors of the PLA2 cascade markedly inhibit LPS-induced TNF-alpha release, supporting a key role of this pathway in the regulation of microglial activation. We also investigated the putative effects of cGMP-elevating agents on blocking microglial activation induced by LPS. Data show that each member of this class of cGMP-elevating compounds that we employed opposed microglial TNF-alpha release, suggesting that strengthening intracellular cGMP signaling mitigates against microglial activation. Taken together, our results suggest novel strategies for reducing microglial activation.

L3 ANSWER 5 OF 8 MEDLINE on STN  
ACCESSION NUMBER: 1999318341 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 10391457  
TITLE: Involvement of axonal phospholipase A2 activity in the outgrowth of adult mouse sensory axons in vitro.  
AUTHOR: Hornfelt M; Ekstrom P A; Edstrom A  
CORPORATE SOURCE: Department of Animal Physiology, Lund University, Sweden.  
SOURCE: Neuroscience, (1999) Vol. 91, No. 4, pp. 1539-47.  
Journal code: 7605074. ISSN: 0306-4522.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199908  
ENTRY DATE: Entered STN: 10 Sep 1999  
Last Updated on STN: 10 Sep 1999  
Entered Medline: 24 Aug 1999

AB The effect on axonal outgrowth of inhibition of phospholipase A2 activity was studied in a recently developed in vitro model, where dorsal root ganglia with attached spinal roots and nerve stumps from young adult mice were cultured in an extracellular matrix material (Matrigel). The phospholipase A2 inhibitors 4-bromophenacyl bromide and oleyloxyethyl phosphorylcholine dose-dependently reduced axonal outgrowth from the sciatic nerve stump. A similar inhibitory effect was seen when only the cut nerve end was exposed to the inhibitors in a compartmental culture system. The local effect of phospholipase A2 inhibition was further investigated on axons established in culture, using time-lapse recording. Exposure to phospholipase A2 inhibitors caused the retraction of filopodia extensions and a reduction in growth cone motility within a few minutes. After removal of inhibition, normal growth cone motility and axonal growth were regained. Nerve cell bodies and axons, in contrast to Schwann cells, showed immunoreactivity after staining with an antiserum against secretory phospholipase A2, and elevated levels of the enzyme could be detected

after culture for 24 h. The immunoreactive protein was of approximately 170,000 molecular weight (phospholipase A2-170) as determined by sodium dodecyl sulphate-polyacrylamide gel electrophoresis and immunoblotting. The localization of phospholipase A2-170 in axons growing into the Matrigel was also demonstrated by use of a whole-mount technique. The results of this study show the importance of continuous phospholipase A2 activity for growth cone motility and axonal outgrowth in the mammalian peripheral nerve, and suggest the involvement of an axonally localized enzyme.

L3 ANSWER 6 OF 8 MEDLINE on STN  
ACCESSION NUMBER: 1999217849 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 10203186  
TITLE: Role of endothelial factors in the specific response of mouse tumour-feeding arterioles to stimulation of 5-HT<sub>1</sub> receptors.  
AUTHOR: Laemmel E; Stucker O; Vicaut E  
CORPORATE SOURCE: Dept de Biophysique et INSERM U141, Hopital F. Widal, Paris, France.  
SOURCE: International journal of radiation biology, (1999 Mar) Vol. 75, No. 3, pp. 365-71.  
Journal code: 8809243. ISSN: 0955-3002.  
PUB. COUNTRY: ENGLAND: United Kingdom  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals; Space Life Sciences  
ENTRY MONTH: 199904  
ENTRY DATE: Entered STN: 11 May 1999  
Last Updated on STN: 11 May 1999  
Entered Medline: 29 Apr 1999

AB PURPOSE: To investigate the possible role of endothelial mediators on the increased vasoconstriction to 5-HT<sub>1</sub> receptor stimulation by the host-modified arterioles feeding a Meth-A tumour implanted in the flank of female Balb/c mice. MATERIALS AND METHODS: Using intravital microscopy, the response to the topical administration of the general 5-HT<sub>1</sub> agonist 5-carboxamidotryptamine maleate (5-CT; 10(-6) M to 10(-4) M) by the tumour-feeding arterioles with the responses of tumour-independent arterioles and those of control arterioles from mice without tumour after antagonization or inhibition of the synthesis of endothelial mediators was compared. RESULTS: The dramatically higher vasoconstriction to 5-CT observed in tumour-feeding arterioles than in tumour-independent or control arterioles still persisted when either nitric oxide synthase, cyclooxygenase, lipoxigenase, or phospholipase A2 were inhibited or when thromboxane A2 or endothelin were antagonized. CONCLUSIONS: It was concluded that the higher reactivity to 5-HT<sub>1</sub> stimulation by tumour-feeding arterioles is not due to changes in endothelial mediator release but probably due to changes affecting arteriolar smooth muscle.

L3 ANSWER 7 OF 8 MEDLINE on STN  
ACCESSION NUMBER: 91183640 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 1901255  
TITLE: Inhibitors of cytochrome P-450 attenuate the myogenic response of dog renal arcuate arteries.  
AUTHOR: Kauser K; Clark J E; Masters B S; Ortiz de Montellano P R; Ma Y H; Harder D R; Roman R J  
CORPORATE SOURCE: Department of Physiology, Medical College of Wisconsin, Milwaukee 53226.  
CONTRACT NUMBER: HL-29587 (NHLBI)  
HL-33833 (NHLBI)  
HL-36279 (NHLBI)  
+  
SOURCE: Circulation research, (1991 Apr) Vol. 68, No. 4, pp. 1154-63.  
Journal code: 0047103. ISSN: 0009-7330.  
PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)  
(RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 199105

ENTRY DATE: Entered STN: 26 May 1991

Last Updated on STN: 3 Feb 1997

Entered Medline: 8 May 1991

AB The role of cytochrome P-450 in the myogenic response of isolated, perfused renal arcuate arteries of dogs to elevations in transmural pressure was examined. The phospholipase A2 inhibitor oleyloxyethylphosphorylcholine (1 and 10 microM) inhibited the greater than threefold increase in active wall tension in these arteries after an elevation in perfusion pressure from 80 to 160 mm Hg. Inhibition of cyclooxygenase activity with indomethacin (1 or 10 microM) had no effect on this response. The cytochrome P-450 inhibitors ketoconazole (10 and 100 microM) and beta-diethyl-aminoethyldiphenylpropylacetate (SKF 525A, 10 and 100 microM) also inhibited the myogenic response. At a pressure of 160 mm Hg, SKF 525A (10 microM) and ketoconazole (100 microM) reduced active wall tension in renal arteries by approximately 70%. Partial inhibition of the myogenic response was obtained after perfusion of the vessels with mechanism-based inhibitors of P-450, 1-aminobenzotriazole (75 microM) and 12-hydroxy-16-heptadecynoic acid (20 microM). The thromboxane receptor antagonist SQ 29,548 (1 or 10 microM) had no effect on the pressure-induced increase in active wall tension in renal arteries. Arachidonic acid (50 microM) constricted isolated perfused renal arteries and potentiated the myogenic response in the presence of indomethacin. This response was completely reversed by ketoconazole (100 microM) or SKF 525A (100 microM). Microsomes (1 mg/ml) prepared from small renal arteries (200-500 microns) and incubated with [1-14C]arachidonic acid (0.5 mu Ci, 50 microM) produced a metabolite that coeluted with 20-hydroxyeicosatetraenoic acid (20-HETE) during reversed-phase high-performance liquid chromatography. The formation of this product was inhibited by both ketoconazole and SKF 525A at concentrations of 10 and 100 microM. These results are consistent with the involvement of the vasoconstrictor 20-HETE and other cytochrome P-450 metabolites of endogenous fatty acids in the myogenic response.

L3 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1985:406131 CAPLUS

DOCUMENT NUMBER: 103:6131

ORIGINAL REFERENCE NO.: 103:1103a,1106a

TITLE: A new efficient and versatile synthesis of alkyl phosphorylcholines

AUTHOR(S): Magolda, R. L.; Johnson, P. R.

CORPORATE SOURCE: Cent. Res. Dev. Dep., E. I. du Pont de Nemours and Co., Wilmington, DE, 19898, USA

SOURCE: Tetrahedron Letters (1985), 26(9), 1167-70

CODEN: TELEAY; ISSN: 0040-4039

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 103:6131


AB Phosphorylcholines ROP(O)(O-)OCH2CH2N+Me3 [R = Me(CH2)n, Me(CH2)7CH:CH(CH2)8, Me(CH2)mS(CH2)3, Me(CH2)7CH:CH(CH2)8S(CH2)3, Me(CH2)mOCH2CH2, Me(CH2)7CH:CH(CH2)8OCH2CH2; m = 15,17; r = 5,7,11,17] were prepared in 35-50% overall yield by treating ROH with POCl3, followed by ethylene glycol and treating the resulting cyclic phosphates with Me3N.

IT 96720-06-8P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 96720-06-8 CAPLUS

CN 3,5,8-Trioxa-4-phosphahexacos-17-en-1-aminium, 4-hydroxy-N,N,N-trimethyl-, inner salt, 4-oxide (9CI) (CA INDEX NAME)



=> file registry

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
15.53	22.96

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-0.80	-0.80

CA SUBSCRIBER PRICE

FILE 'REGISTRY' ENTERED AT 16:52:52 ON 25 JAN 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 24 JAN 2008 HIGHEST RN 1000773-19-2

DICTIONARY FILE UPDATES: 24 JAN 2008 HIGHEST RN 1000773-19-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=> E "OOPC"/CN 25

E1	1	OOLONGTHEANIN-3'-O-GALLATE/CN
E2	1	OOMYCIN A/CN
E3	0 -->	OOPC/CN
E4	1	OOPG 1000/CN
E5	1	OOPG 1002/CN
E6	1	OOPASM SPECIFIC PROTEIN (MUS MUSCULUS STRAIN NIH/SWISS GENE OP1)/CN
E7	1	OOPODIN/CN
E8	1	OOPODIN, 11,13-DIDEHYDRO-/CN
E9	1	OOPORPHYRIN/CN
E10	1	OORA SUBUNIT OF 2-OXOGLUTARATE:ACCEPTOR OXIDOREDUCTASE (CAMPYLOBACTER JEJUNI STRAIN NCTC 11168 GENE OORA)/CN
E11	1	OORA SUBUNIT OF 2-OXOGLUTARATE:ACCEPTOR OXIDOREDUCTASE (HELICOBACTER HEPATICUS STRAIN ATCC51449 GENE OORA)/CN
E12	1	OORAKKU APO 101/CN
E13	1	OORAKKU APO 101-HITALOID 3083-70B-MDI COPOLYMER/CN
E14	1	OORAKKU APO 101-HITALOID 3083-70B-MILLIONATE MR 200 COPOLYMER/CN
E15	1	OORAKKU APO 301/CN
E16	1	OORB SUBUNIT OF 2-OXOGLUTARATE:ACCEPTOR OXIDOREDUCTASE (CAMPYLOBACTER JEJUNI STRAIN NCTC 11168 GENE OORB)/CN

E17 1 OORB SUBUNIT OF 2-OXOGLUTARATE:ACCEPTOR OXIDOREDUCTASE  
 (HELICOBACTER HEPATICUS STRAIN ATCC51449 GENE OORB)/CN  
 E18 1 OORC SUBUNIT OF 2-OXOGLUTARATE:ACCEPTOR OXIDOREDUCTASE  
 (CAMPYLOBACTER JEJUNI STRAIN NCTC 11168 GENE OORC)/CN  
 E19 1 OORC SUBUNIT OF 2-OXOGLUTARATE:ACCEPTOR OXIDOREDUCTASE  
 (HELICOBACTER HEPATICUS STRAIN ATCC51449 GENE OORC)/CN  
 E20 1 OORD SUBUNIT OF 2-OXOGLUTARATE:ACCEPTOR OXIDOREDUCTASE  
 (CAMPYLOBACTER JEJUNI STRAIN NCTC 11168 GENE OORD)/CN  
 E21 1 OORD SUBUNIT OF 2-OXOGLUTARATE:ACCEPTOR OXIDOREDUCTASE  
 (HELICOBACTER HEPATICUS STRAIN ATCC51449 GENE OORD)/CN  
 E22 1 OORP (ONCORHYNCHUS MYKISS OOCYTE)/CN  
 E23 1 OOSPGLYCOL/CN  
 E24 1 OOSPOALDEHYDE/CN  
 E25 1 OOSPOALDEHYDE, (2,4-DINITROPHENYL)HYDRAZONE/CN

=> d his

(FILE 'HOME' ENTERED AT 16:45:32 ON 25 JAN 2008)

FILE 'REGISTRY' ENTERED AT 16:45:44 ON 25 JAN 2008

E "OLEYLOXYETHYLPHOSPHOCHOLINE"/CN 25

E "OOEPC"/CN 25

L1 1 S 96720-06-8/RN

L2 1 S L1

FILE 'MEDLINE, CAPLUS, WPIDS, USPATFULL' ENTERED AT 16:49:58 ON 25 JAN 2008

L3 8 S L1

FILE 'REGISTRY' ENTERED AT 16:52:52 ON 25 JAN 2008

E "OOPC"/CN 25

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	11.96	34.92
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-0.80

STN INTERNATIONAL LOGOFF AT 17:08:19 ON 25 JAN 2008